

**Supplemental Table 7. Changes in the parameters at the first visit before 2012, visit in 2012, and the last health check visit after 2012 in the participants with CKD Stage G1 in 2012**

	number of data	First Visit			2012			Last Visit			P
		mean	±	SD	mean	±	SD	mean	±	SD	
<b>BMI (Kg/m<sup>2</sup>)</b>	36	<b>22.8</b>	±	3.7	<b>22.7</b>	±	3.9	<b>22.5</b>	±	3.9	0.233
<b>Waist circumference (cm)</b>	35	<b>84.8</b>	±	9.0	<b>83.7</b>	±	9.1	<b>83.3</b>	±	9.6	0.146
<b>SBP (mmHg)</b>	36	<b>126</b>	±	17	<b>124</b>	±	13	<b>125</b>	±	15	0.683
<b>DBP (mmHg)</b>	36	<b>76</b>	±	11	<b>75</b>	±	11	<b>75</b>	±	10	0.792
<b>TG (mg/dl)</b>	36	<b>121</b>	±	87	<b>111</b>	±	71	<b>115</b>	±	80	0.550
<b>HDL-C (mg/dl)</b>	36	<b>63</b>	±	14	<b>65</b>	±	16	<b>66</b>	±	16	0.108
<b>LDL-C (mg/dl)</b>	36	<b>126</b>	±	33	<b>121</b>	±	35	<b>124</b>	±	28	0.647
<b>FBS (mg/dl)</b>	31	<b>98</b>	±	17	<b>101<sup>c</sup></b>	±	18	<b>96<sup>b</sup></b>	±	17	0.007
<b>HbA1c (%)</b>	36	<b>5.3<sup>c</sup></b>	±	0.6	<b>5.3<sup>c</sup></b>	±	0.7	<b>5.7<sup>a,b</sup></b>	±	0.7	0.000
<b>eGFR (ml/min/1.73m<sup>2</sup>)</b>	36	<b>91.7<sup>c</sup></b>	±	14.1	<b>100.0<sup>c</sup></b>	±	16.2	<b>84.1<sup>a,b</sup></b>	±	13.3	0.000
<b>Uric acid (mg/dl)</b>	36	<b>4.6</b>	±	1.1	<b>4.6</b>	±	0.9	<b>4.8</b>	±	1.0	0.237
<b>Urinary protein (0-3)</b>	35	<b>0.200<sup>b</sup></b>	±	0.424	<b>0.629<sup>a,c</sup></b>	±	0.634	<b>0.243<sup>b</sup></b>	±	0.547	0.000
<b>Urinary occult blood (0-3)</b>	32	<b>1.047<sup>c</sup></b>	±	1.297	<b>1.359<sup>c</sup></b>	±	1.079	<b>0.469<sup>a,b</sup></b>	±	0.851	0.002
<b>Urine sugar (0-3)</b>	35	<b>0.000</b>	±	0.000	<b>0.029</b>	±	0.169	<b>0.000</b>	±	0.000	0.368

Abbreviations are: SD; standard deviation, BMI; body mass index, SBP; systolic blood pressure, DBP; diastolic blood pressure, TG; triglyceride, HDL-C; high density lipoprotein cholesterol, LDL-C; low density lipoprotein cholesterol, FBS; fast blood sugar, Hb; hemoglobin

<sup>a</sup> Significantly differ compared to the first visit

<sup>b</sup> Significantly differ compared to visit in 2012

<sup>c</sup> Significantly differ compared to the last visit

As for the comparisons of urinary findings between two points among three points, p<0.0167 was defined to be statistically significant.